

Supplementary information, Fig. S3. In this figure, cancer cells were treated with DM- α KG (15 mM) for 24 hours to assess DR6 oxidation and pyroptotic features (including morphology, GSDMC cleavage, and LDH release), unless specially indicated otherwise.

(a) Inhibitors, Genistein or M β CD, showed no effect on DM- α KG-induced DR6 oxidation. HeLa cells were pretreated with Genistein and M β CD for 2 hours.

(b) The knockdown efficiency of CLTA, CLTB, CLAC, DNM1 and DNM2 in HeLa cells as determined by RT-qPCR.

(c, d, e) Effect of CLTs on DM- α KG-induced caspase-8 activation (d), pyroptotic morphology (d), GSDMC cleavage and LDH release (e). CLTA, CLTB, and CLAC had first been knocked down in HeLa cells.

(f, g, h) Effect of DNMs on DM- α KG-induced pyroptotic morphology (f), caspase-8 activation (g), GSDMC cleavage and LDH release (h). DNM1 and DNM2 had first been knocked down in the cells.


(i) Genistein or M β CD impaired DM- α KG-induced GSDMC cleavage in SGC-7901 (top) and B16 (bottom) cells.

Tubulin was used to determine the amount of loading proteins. All data are presented as the mean \pm SEM of two or three independent experiments. *** $p < 0.001$. The data


were analyzed using two-way ANOVA followed by the Bonferroni test.

Supplementary information, Figure S3


a








b



c



d**e**



i

